



Note. Dotted lines indicate confidence intervals; solid lines indicate regression model results.

FIGURE 1—Acute myocardial infarction mortality rates in Jefferson County, TX, where effective tobacco control activities were carried out in 2000–2005, and other Texas counties for 1996–2000 and 2001–2005.

However, the promise shown by this pilot project was not fulfilled in Texas. In 2006, tobacco programs at the Texas Department of State Health Services were reorganized, and their funding was reduced. Support for activities in Jefferson County was withdrawn. ■

About the Authors

Alfred L. McAlister, Philip Huang, and Ronald B. Harrist are with the University of Texas School of Public Health, Austin Regional Campus, Austin. Philip Huang is also with the Austin/Travis County Health and Human Services Department, Austin. Amelie G. Ramirez is with the University of Texas Health Science Center, San Antonio. At the time of the study, Vincent P. Fonseca was with the Texas Department of State Health Services, Austin.

Correspondence should be sent to Alfred L. McAlister, University of Texas School of Public Health, Austin Regional Campus, 1616 Guadalupe St, Suite 6.300, Austin TX 78701 (e-mail: Alfred.L.McAlister@uth.tmc.edu). Reprints can be ordered at <http://www.ajph.org> by clicking the "Reprints/Eprints" link.

This article was accepted March 17, 2010.

Contributors

A.L. McAlister and P. Huang were the primary writers. A.G. Ramirez wrote the description of intervention activities and edited the Discussion section. R.B. Harrist conducted analyses, prepared the figure, and wrote most of the Results section. V.P. Fonseca collected the data and wrote parts of the Introduction and Results sections.

Acknowledgments

This research was supported in part by a grant from the National Institutes of Health (R01-CA-86295).

Human Participant Protection

No protocol approval was required because only publicly available data were used.

References

- McAlister A, Morrison TC, Hu S, et al. Media and community campaign effects on adult tobacco use in Texas. *J Health Commun.* 2004;9(2):95–109.
- McAlister AL, Rabinus V, Geiger A, Glynn TJ, Huang P, Todd R. Telephone assistance for smoking cessation: one year cost effectiveness estimations. *Tob Control.* 2004; 13(1):85–86.
- McAlister AL, Huang P, Ramirez AG. Settlement-funded tobacco control in Texas: 2000–2004 pilot project effects on cigarette smoking. *Public Health Rep.* 2006;121(3):235–238.
- International Classification of Diseases, 10th Revision. Geneva, Switzerland: World Health Organization; 1992.
- Anderson RN, Rosenberg HM. Age standardization of death rates: implementation of the year 2000 standard. *Natl Vital Stat Rep.* 1998;47(3):1–16, 20.
- Schroeder SA. Public smoking bans are good for the heart. *J Am Coll Cardiol.* 2009;54(14):1256–1257.
- Sargent RP, Shepard RM, Glantz SA. Reduced incidence of admissions for myocardial infarction

associated with public smoking ban: before and after study. *BMJ.* 2004;328(7446):977–980.

8. Vartiainen E, Puska P, Pekkanen P, Toumilehto J, Jousilahti P. Changes in risk factors explain changes in mortality from ischaemic heart disease in Finland. *BMJ.* 1994;309(6946):23–27.

Adding Sexual Orientation Questions to Statewide Public Health Surveillance: New Mexico's Experience

Nicole A. VanKim, MPH, James L. Padilla, MS, Joseph G.L. Lee, MPH, CPH, and Adam O. Goldstein, MD, MPH

We examined refusal rates for sensitive demographic questions to determine whether questions on sexual orientation are too sensitive for routine use on public health surveys. We compared the percentage of active refusals in New Mexico for a sexual orientation question and 6 other sensitive demographic questions. In 2007 and 2008, refusal rates for sexual orientation questions were similar to rates for questions on race/ethnicity and weight and significantly lower than rates for questions on household income. Perceptions that sexual orientation is too controversial a topic to be included on state surveys may be unfounded. (*Am J Public Health.* 2010;100:2392–2396. doi:10.2105/AJPH.2009.186270)

Lesbian, gay, bisexual, and transgender (LGBT) populations have clear disparities in cigarette smoking,^{1,2} suicidal ideation,³ violent victimization,^{4–6} and sexually transmitted infections⁷ compared with the general population. Yet demographic questions about sexual orientation (i.e., questions about identity, attraction, or behavior), in addition to those on gender identity,

are often not included in routine health surveys.^{8–11}

The Behavioral Risk Factor Surveillance System (BRFSS) survey, Adult Tobacco Survey (ATS), Youth Risk Behavior Survey, and Youth Tobacco Survey make up the backbone of state health surveillance in the United States, providing benchmarks, informing interventions, and allowing for comparisons between states. In 1998, the Centers for Disease Control and Prevention (CDC) chose not to include a question on sexual orientation in the Youth Risk Behavior Survey, leaving the decision to individual states,¹² which collaboratively administer a core survey and state-added modules. This decision has remained in place and covers, in practice, other CDC-run, state-administered surveys such as the BRFSS, ATS, and Youth Tobacco Survey. Over the subsequent 12 years, approximately 13 states have collected information on sexual orientation in the BRFSS¹³; California does so by using the separate California Health Interview Survey (<http://www.chis.ucla.edu>). Even fewer states collect such data on other statewide surveys.¹⁴

No research has examined why states do not include such questions on their surveys. Several possible explanations exist; public health practitioners may (1) feel that the evidence of health disparities is not enough to warrant inclusion of such questions in state surveys, (2) perceive a lack of expertise and capacity to develop and include questions on sexual orientation and gender identity, or (3) believe sexual orientation questions will yield a high nonresponse rate or cause early survey termination because of respondents' discomfort with the topic. A fourth possibility is that decision-makers may not support inclusion of such questions because of political or personal biases.

Strong evidence and resources are available to address the first⁹ and second concerns; technical assistance is available to enhance capacity and cultural competency through the National LGBT Tobacco Control Network (<http://www.lgbttobacco.org>), and the University of California, Los Angeles's (UCLA's) Williams Institute recently released a best-practices guide to asking questions on sexual orientation.¹⁵ We address the third concern—the belief that sexual orientation questions may be too

sensitive for use in public surveys—by using New Mexico's experience in including such questions in 2 statewide population-based surveys.

Addressing the last concern, regarding political and personal bias, is outside the scope of this report.

Early research on sexual orientation questions and refusal rates came from surveys of women and health care professionals^{16–18} and, recently, from surveys in the Pacific Northwest¹⁹ and Massachusetts.²⁰ Using 2003 Oregon and Washington BRFSS data, Dilley et al. reported that 3% of those surveyed reported “do not know” or refused to answer sexual orientation questions.²¹ Dilley et al. reported refusal rates as low as 1.2% to 1.6% in combined 2003–2006 Washington State BRFSS data.¹⁹ In combined 2001–2006 Massachusetts BRFSS data, 3.6% of respondents refused to answer questions on sexual orientation.²² We identified no published research that reported refusal rates for sexual orientation questions by year that showed possible trends or compared refusal rates for sexual orientation questions with refusal rates for other sensitive questions. Although survey researchers have established that inquiries about household income are one of the most sensitive—and therefore most often refused—questions,²³ little research has directly addressed concerns that sexual orientation questions may be too sensitive for the public in states outside of the Pacific Northwest and New England. We examined the sensitivity of questions on sexual orientation and other demographic characteristics by using results from New Mexico health surveys between 2003 and 2008.

METHODS

In 2003, the New Mexico Department of Health's Tobacco Use Prevention and Control (TUPAC) Program identified lesbian, gay, and bisexual people as a priority population in addressing tobacco-related disparities. TUPAC then successfully advocated for inclusion of the following question on the New Mexico ATS: “Do you consider yourself to be—?” Possible responses included “heterosexual or straight,” “homosexual (gay or lesbian),” “bisexual,” “other,” and “don't know/not sure,” or the respondent could refuse to answer. The 2003 ATS became the first statewide population-based health

surveillance system to assess sexual orientation in New Mexico.

The question's wording was adapted from the CDC's National Health and Nutrition Examination Survey²⁴ and was placed at the end of the ATS demographics section, which was prefaced with an explanation that there might be differences based on race, ethnicity, household income, sexual orientation, and education level. In a pilot study, telephone surveyors from the New Mexico Department of Health Survey Section asked the sexual orientation question to 25 to 30 randomly selected residents, documenting in turn the questions asked by residents and their difficulty with the wording of the question.

Using the experience of the New Mexico ATS and the results from the pilot study, the 2005 and 2006 New Mexico BRFSS included a state-added question on sexual orientation. Unlike the question on the New Mexico ATS, this BRFSS question did not have a preface. Instead, the following transition statement was used in the BRFSS: “Now I'm going to ask you a question about sexual orientation.” Surveyors were trained in administering the question and were provided talking points and detailed background information for use in responding to callers who may have had concerns about the purpose of the question. Surveyors reported that some older respondents seemed confused when asked the sexual orientation question. A significantly higher percentage of adults aged 65 years and older responded “don't know” (data not shown). This question was therefore administered only to respondents aged 18 to 64 years on the 2007 and 2008 New Mexico BRFSS.

Surveys

Both the New Mexico ATS and New Mexico BRFSS are ongoing, population-based, random-digit-dialed, landline telephone surveys of English- or Spanish-speaking noninstitutionalized adults who live in a household in New Mexico. The New Mexico ATS contains questions assessing tobacco-related behaviors and attitudes. The New Mexico BRFSS has questions assessing various health characteristics, including risk factors, preventive factors, and chronic diseases. Both surveys include a demographics section containing questions on

race, ethnicity, age, gender, education level, and household income. On the New Mexico ATS, sexual orientation is asked as part of the demographics section. On the New Mexico BRFSS, sexual orientation is asked in a state-added question and is thus located toward the end of the survey.

Additional in-depth documentation on the BRFSS is publicly available online (<http://www.cdc.gov/brfss>). The ATS uses BRFSS methodology, albeit with different content and sample size.

Analysis

For both the New Mexico ATS and the New Mexico BRFSS, responses receive 1 of 4 codes: direct response (e.g., straight, gay, lesbian, bisexual, other), missing, “refused,” or “don’t know.” Missing responses indicate that the respondent did not receive the question, often because of skip patterns (e.g., surveyors did not ask respondents aged 65 years or older the sexual orientation question in the 2007 and 2008 New Mexico BRFSS) or because of early termination of the survey. Missing values were not included in the analyses. Responses coded as “refused” indicate that the respondent actively refused to provide an answer to the question. Responses coded as “don’t know” indicate that the respondent did not select one of the survey responses but also did not

actively refuse to provide an answer. For this study, we compared the percentages of refusals for the sexual orientation question and for 6 other demographic questions: on household income, education level, race/ethnicity, Hispanic origin, age, and body weight.

Since both the New Mexico ATS and the New Mexico BRFSS are landline telephone-based surveys, they are subject to sampling bias. To help adjust for bias, responses are weighted on each respondent’s probability of selection and stratified to the gender, age, and geographic distribution of New Mexico. Both surveys require completion through the demographics section for the record to be included in the data set.

We used only data sets that assessed sexual orientation. Thus, for the New Mexico BRFSS, we used data sets for 2005 ($n=5585$), 2006 ($n=6581$), 2007 ($n=6606$), and 2008 ($n=6227$). For the New Mexico ATS, we used data sets for 2003 ($n=2503$) and 2006 ($n=2551$). Annual response rates ranged from 55.5% to 58.9% on the New Mexico BRFSS²⁵ and from 45.9% to 48.0% on the New Mexico ATS.²⁶ Nonoverlapping confidence intervals (CIs) indicate statistical significance at $P<.05$. To account for weighting, we conducted statistical analyses by using SVY commands in Stata version 10 (Stata Corp LP, College Station, TX).

RESULTS

Table 1 presents sample sizes and the weighted percentages (with 95% CIs) of respondents who refused to answer questions on sexual orientation and on selected demographic characteristics in the New Mexico ATS and the New Mexico BRFSS. The question on household income had a significantly higher rate of refusal than did questions on other demographic characteristics, including sexual orientation, for all years of the surveys. Rates of refusal for the household income question ranged from 4.23% to 6.11%, whereas refusal rates for the sexual orientation question ranged from 0.80% to 2.60% and refusal rates for other questions on demographic characteristics ranged from 0.05% to 1.37%. Refusal rates for the question on sexual orientation were significantly higher than for questions on all demographic characteristics except household income on the 2003 and 2006 New Mexico ATS. There were no significant differences in refusal rates for questions on sexual orientation and body weight in the 2005 and 2006 New Mexico BRFSS.

In the 2007 New Mexico BRFSS, the percentage of respondents who refused to answer the question on sexual orientation (0.80%; 95% CI=0.53, 1.20) was significantly higher than the percentage who refused to answer

TABLE 1—Number and Weighted Percentage of Respondents Who Actively Refused to Answer Questions on Sexual Orientation and Selected Demographic Characteristics: New Mexico Adult Tobacco Survey (ATS) and Behavioral Risk Factor Surveillance System (BRFSS), 2003–2008

	2003 ATS (n=2503)		2005 BRFSS (n=5585)		2006 ATS (n=2551)		2006 BRFSS (n=6581)		2007 BRFSS (n=6606)		2008 BRFSS (n=6227)	
	No. of Refusals	Weighted % (95% CI)	No. of Refusals	Weighted % (95% CI)	No. of Refusals	Weighted % (95% CI)	No. of Refusals	Weighted % (95% CI)	No. of Refusals	Weighted % (95% CI)	No. of Refusals	Weighted % (95% CI)
Sexual orientation	52	1.85 (1.32, 2.58)	94	1.18 (0.92, 1.50)	98	2.60 (2.08, 3.25)	106	1.09 (0.86, 1.38)	42	0.80 ^a (0.53, 1.20)	55	1.13 ^a (0.75, 1.71)
Age	22	0.56 (0.35, 0.90)	12	0.21 (0.11, 0.41)	20	0.58 (0.35, 0.96)	39	0.49 (0.33, 0.73)	25	0.31 (0.19, 0.49)	16	0.25 (0.14, 0.46)
Hispanic origin	8	0.23 (0.10, 0.48)	16	0.28 (0.15, 0.51)	10	0.31 (0.15, 0.60)	30	0.39 (0.26, 0.59)	16	0.24 (0.14, 0.43)	20	0.44 (0.19, 0.99)
Race/Ethnicity	21	0.74 (0.46, 1.18)	33	0.54 (0.35, 0.83)	21	0.54 (0.33, 0.88)	55	0.95 (0.66, 1.40)	36	0.49 (0.34, 0.72)	35	0.70 (0.38, 1.28)
Education level	12	0.46 (0.24, 0.87)	16	0.19 (0.11, 0.34)	11	0.28 (0.15, 0.52)	15	0.14 (0.08, 0.27)	8	0.08 (0.03, 0.22)	6	0.05 (0.02, 0.13)
Household income	172	6.00 (5.05, 7.11)	345	5.15 (4.54, 5.83)	177	6.11 (5.12, 7.29)	356	5.02 (4.41, 5.71)	310	4.23 (3.67, 4.88)	322	4.33 (3.66, 5.13)
Body weight	NA ^b	NA ^b	96	1.37 (1.07, 1.75)	NA ^b	NA ^b	94	1.18 (0.88, 1.56)	67	0.91 (0.66, 1.24)	70	0.83 (0.60, 1.14)

Note. CI=confidence interval; NA=not asked. “Refusal” includes only responses coded as “refused.” Percentages are weighted for each respondent’s probability of selection.

^aIncludes only adults aged 18 to 64 years (2007: $n=4840$; 2008: $n=4455$); adults aged 65 years and older were coded as missing and therefore not included in the denominator.

^bQuestion not asked on survey.

questions on age (0.31%; 95% CI=0.19, 0.49), Hispanic origin (0.24%; 95% CI=0.14, 0.43), and education level (0.08%; 95% CI=0.03, 0.22). No significant differences were found in refusal rates for questions on sexual orientation, race/ethnicity (0.49%; 95% CI=0.34, 0.72), and body weight (0.91%; 95% CI=0.66, 1.24) in 2007. In the 2008 New Mexico BRFSS, refusal rates for questions on sexual orientation (1.13%; 95% CI=0.75, 1.71) were significantly higher than for questions on age (0.25%; 95% CI=0.14, 0.46) and education level (0.05%; 95% CI=0.02, 0.13). There were no significant differences between refusal rates for questions on sexual orientation, Hispanic origin (0.44%; 95% CI=0.19, 0.99), race/ethnicity (0.70%; 95% CI=0.38, 1.28), and body weight (0.83%; 95% CI=0.60, 1.14) in 2008.

DISCUSSION

In New Mexico from 2003–2008, refusal rates for a question on sexual orientation were low (0.80%–2.60%). These rates were notably lower than refusal rates for a question on household income (4.23%–6.11%, compared with a refusal rate as high as 12% in the combined Washington–Oregon 2003 BRFSS²¹) and similar to refusal rates for a question on body weight. The lower refusal rates for sexual orientation questions are similar to findings from Massachusetts's 2001–2006 BRFSS (3.6%),²² Washington's 2003–2006 BRFSS (1.2%–1.6%),¹⁹ and combined Washington–Oregon BRFSS data from 2003 (3%).²¹

Our findings in New Mexico corroborate findings from New England and the Pacific Northwest, and our findings do so across 4 years of survey data from 2 separate statewide surveys. Additionally, because New Mexico is a rural border state with a population that is "minority majority" (i.e., minority groups outnumber Whites), our findings suggest that states demographically quite different from states in New England and the Pacific Northwest can successfully improve surveillance systems and maintain data quality by adding a sexual orientation question.

To continue improving their quality of data collection, the 2009 New Mexico ATS and New Mexico BRFSS used a revised sexual

orientation question that assesses both sexual orientation and gender identity: "Do you consider yourself to be one or more of the following—? Possible responses include "straight," "gay or lesbian," "bisexual," "transgender," "other," "don't know/not sure," or the respondent could refuse to answer. The revised question, which was developed with cognitive interviewing techniques (i.e., in-depth interviews with survey respondents to identify thought processes behind how questions were answered), more accurately captures those constructs among adults of all ages.^{27,28}

The availability of sexual orientation data is crucial for interventions against existing health disparities.¹¹ States should recognize that, as refusal rates are similar for questions on sexual orientation and body weight, sexual orientation is not too sensitive a subject for use in public surveys. Additionally, with only a quarter of states choosing to collect data on sexual orientation,¹³ the CDC should revisit earlier decisions to make such questions optional on the BRFSS. ■

About the Authors

Nicole A. VanKim and James L. Padilla are with the Chronic Disease Prevention and Control Bureau, New Mexico Department of Health, Albuquerque. Joseph G. L. Lee and Adam O. Goldstein are with the Tobacco Prevention and Evaluation Program, Department of Family Medicine, University of North Carolina School of Medicine, Chapel Hill.

Correspondence should be sent to James L. Padilla, 5301 Central Ave NE, Suite 800, Albuquerque, NM 87108 (e-mail: james.padilla@state.nm.us). Reprints can be ordered at <http://www.ajph.org> by clicking the "Reprints/Eprints" link.

This brief was accepted March 21, 2010.

Contributors

N.A. VanKim performed all data analyses, drafted the Methods and Results sections of the article, and provided feedback on all drafts of the article. J.L. Padilla assisted in data analyses and data interpretation and provided critical feedback on all drafts of the article. J.G.L. Lee originated the study, drafted the article's introduction and conclusions, and provided feedback on all drafts of the article. A.O. Goldstein provided guidance and critical feedback on all drafts of the article.

Acknowledgments

This report was supported in part by an appointment to the Applied Epidemiology Fellowship Program, administered by the Council of State and Territorial Epidemiologists and funded by Centers for Disease Control and Prevention (Cooperative Agreement U60/CCU007277).

We thank Wayne Honey and Annie Hickman of the New Mexico Department of Health Survey Section and Anna McCullough and Kathryn D. Kramer of the University of North Carolina School of Medicine for helpful comments on the draft.

Note. The opinions expressed herein are those of the authors and do not necessarily represent the policies or opinions of the funders.

Human Participant Protection

No institutional review board approval was needed for this study because data were collected anonymously from a public health surveillance system in which adults voluntarily consented to telephone interviews.

References

1. Gruskin EP, Greenwood GL, Matevia M, Pollack LM, Bye LL. Disparities in smoking between the lesbian, gay, and bisexual population and the general population in California. *Am J Public Health*. 2007;97(8):1496–1502.
2. Lee JG, Griffin GK, Melvin CL. Tobacco use among sexual minorities in the USA, 1987 to May 2007: a systematic review. *Tob Control*. 2009;18(4):275–282.
3. King M, Semlyen J, Tai SS, et al. A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry*. 2008; 8:70.
4. Balsam KF, Rothblum ED, Beauchaine TP. Victimization over the life span: a comparison of lesbian, gay, bisexual, and heterosexual siblings. *J Consult Clin Psychol*. 2005;73(3):477–487.
5. Bontempo DE, D'Augelli AR. Effects of at-school victimization and sexual orientation on lesbian, gay, or bisexual youths' health risk behavior. *J Adolesc Health*. 2002;30(5):364–374.
6. Udry JR, Chantala K. Risk assessment of adolescents with same-sex relationships. *J Adolesc Health*. 2002;31(1):84–92.
7. Valdiserri RO. sexually transmitted infections among gay and bisexual men. In: Wolitski RJ, Stall R, Valdiserri RO, eds. *Unequal Opportunity: Health Disparities Affecting Gay and Bisexual Men in the United States*. New York, NY: Oxford University Press; 2008: 159–193.
8. Sell RL, Becker JB. Sexual orientation data collection and progress toward Healthy People 2010. *Am J Public Health*. 2001;91(6):876–882.
9. Mayer KH, Bradford JB, Makadon HJ, Stall R, Goldhammer H, Landers S. Sexual and gender minority health: what we know and what needs to be done. *Am J Public Health*. 2008;98(6):989–995.
10. Sell RL, Dunn PM. Inclusion of lesbian, gay, bisexual and transgender people in tobacco use-related surveillance and epidemiological research. *J LGBT Health Res*. 2008;4(1):27–42.
11. Gold M, Dodd AH, Neuman M. Availability of data to measure disparities in leading health indicators at the state and local levels. *J Public Health Manag Pract*. 2008;14(suppl):S36–S44.
12. CDC to let states decide whether to ask teens about sex. *AIDS Policy Law*. 1998;13(14):7.
13. Lee JG. Social ecology of tobacco surveillance data for sexual and gender minority populations. *Nicotine Tob Res*. 2009;11(7):908–909.

14. Sell RL. Gay data. 2008. Available at: <http://gaydata.org>. Accessed September 15, 2009.
15. Badgett L, Goldberg N. Best practices for asking questions about sexual orientation on surveys. Williams Institute Sexual Minority Assessment Research Team, Los Angeles, CA, 2009. Available at: http://www.law.ucla.edu/williamsinstitute/pdf/SMART_FINAL_Nov09.pdf. Accessed September 15, 2009.
16. Bradford JB, Honnold JA, Ryan C. Disclosure of sexual orientation in survey research on women. *J Gay Lesbian Med Assoc*. 1997;1(3):169–177.
17. Case P, Austin SB, Hunter DJ, et al. Disclosure of sexual orientation and behavior in the Nurses' Health Study II: results from a pilot study. *J Homosex*. 2006; 51(1):13–31.
18. Brogan D, Frank E, Elon L, O'Hanlan KA. Methodologic concerns in defining lesbian for health research. *Epidemiology*. 2001;12(1):109–113.
19. Dilley JA, Simmons KW, Boysun MJ, Pizacani BA, Stark MJ. Demonstrating the importance and feasibility of including sexual orientation in public health surveys: health disparities in the Pacific Northwest. *Am J Public Health*. 2010;100(3):460–467.
20. Keyes SM, Rothman EF, Zhang Z. Sexual orientation and sexual behavior: results from the Massachusetts Behavioral Risk Factor Surveillance System, 2002–2006. *J LGBT Health Res*. 2007;3(3):1–10.
21. Dilley JA, Maher JE, Boysun MJ, et al. Response letter to: Tang H, Greenwood GL, Cowling DW, Lloyd JC, Roeseler AG, Bal DG. Cigarette smoking among lesbians, gays, and bisexuals: how serious a problem? *Cancer Causes Control*. 2005;16(9):1133–1134.
22. Conron KJ, Mimiaga MJ, Landers SJ. *A Health Profile of Massachusetts Adults by Sexual Orientation Identity: Results From the 2001–2006 Behavioral Risk Factor Surveillance System Surveys*. Boston: Massachusetts Dept of Public Health; 2008.
23. Tourangeau R, Yan T. Sensitive questions in surveys. *Psychol Bull*. 2007;133(5):859–883.
24. Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey. Available at: <http://www.cdc.gov/nchs/nhanes.htm>. Accessed September 23, 2010.
25. Centers for Disease Control and Prevention. BRFSS annual survey data: summary data quality reports. 2009. Available at: http://www.cdc.gov/BRFSS/technical_infodata/quality.htm. Accessed January 12, 2010.
26. Mariolis P. *Guidelines for Conducting General Population State Adult Tobacco Telephone Surveys*. Atlanta, GA: Centers for Disease Control and Prevention; 2007.
27. Clark MA, Armstrong G, Bonacore L. Measuring sexual orientation and gender expression among middle-aged and older women in a cancer screening study. *J Cancer Educ*. 2005;20(2):108–112.
28. Scout, Senseman SE. Cognitive testing of an LGBT surveillance question. 2008. Available at: http://www.lgbttobacco.org/files/2008_09_07_cog_testing_article_briefing.doc. Accessed September 15, 2009.